

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA337164

## EBI3 Mouse Monoclonal Antibody [Clone ID: 42N1G4]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	42N1G4
Applications:	FC, WB
Recommended Dilution:	Flow Cytometry: 0.1-1ug/10^6cells, Western Blot: 2-5ug/ml~
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2b, kappa
Clonality:	Monoclonal
Immunogen:	Full-length recombinant human EBI3 protein full length recombinant human EBI3 (cat # 100- 162)
Formulation:	PBS containing 0.05% BSA, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at - 20C long term. Avoid freeze-thaw cycles.
Concentration:	lot specific
Purification:	Protein G purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	Epstein-Barr virus induced 3
Database Link:	<u>NP_005746</u> <u>Entrez Gene 10148 Human</u> <u>Q14213</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** EBI3 Mouse Monoclonal Antibody [Clone ID: 42N1G4] – TA337164

Background:EBI3 (Epstein-Barr virus Induced-3) is a secreted glycoprotein of the hematopoietin receptor<br/>family. It plays a critical regulatory role in the induction of Th2-type immune responses and<br/>the development of Th2-mediated tissue inflammation in vivo, which may be mediated<br/>through the control of iNkT cell function. EBI3 dimerizes with p28 and p35 subunits of IL-12 to<br/>form new proteins IL-27 and IL-35, respectively . IL-27 is an early product of activated antigen<br/>presenting cell that is produced upon TLR ligation. It negatively regulates Th17 cell<br/>differentiation. EBI3 is widely expressed and its expression in dendritic cell is transcriptionally<br/>regulated by TLR signaling via MyD88 and NF-kappaB during innate immune responses<br/>preceding cytokine driven Th cell development. EBI3 signal inhibits delayed-type<br/>hypersensitivity responses by suppressing IL-17 production and inducing IL-10<br/>hyperproduction. EBI3 may play a novel role in controlling tumor metastasis via lung CD8+ T<br/>cells, and its deficiency is associated with a diminished production of Th2 cytokines which are<br/>known to regulate allergic airway inflammation in asthma. Targeted deletion of EBI3 protects<br/>mice from lung metastasis


Synonyms: IL-27B; IL27B

Protein Families:

L-27B; IL27B


Druggable Genome, Secreted Protein

#### **Product images:**



Western Blot: IL-27/IL-35 EBI3 Subunit Antibody (42N1G4) TA337164 - Analysis using EBI3 antibody. Human EBI3 recombinant protein (A) and human IL-35 recombinant protein (B) probed with EBI3 antibody at 5 ug/ml. Goat anti-mouse Ig HRP secondary antibody and PicoTect ECL substrate solution were used for this test.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Flow Cytometry: IL-27/IL-35 EBI3 Subunit Antibody (42N1G4) TA337164 - Intracellular analysis of EBI3 in transfected cells using EBI3 antibody at 0.2 ug/10<sup>6</sup> cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US