

# **Product datasheet for TA337702**

### **KEAP1 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications:IHC, WBRecommended Dilution:WB, IHCReactivity:HumanHost:RabbitIsotype:IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-KEAP1 antibody: synthetic peptide directed towards the C terminal

of human KEAP1. Synthetic peptide located within the following region: TWTFVAPMKHRRSALGITVHQGRIYVLGGYDGHTFLDSVECYDPDTDTWS

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Protein A purified

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 70 kDa

**Gene Name:** kelch like ECH associated protein 1

Database Link: NP 987096

Entrez Gene 9817 Human

Q14145



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: KEAP1 contains KELCH-1 like domains, as well as a BTB/POZ domain. Kelch-like ECH-

associated protein 1 interacts with NF-E2-related factor 2 in a redox-sensitive manner and the dissociation of the proteins in the cytoplasm is followed by transportation of NF-E2-related factor 2 to the nucleus. This interaction results in the expression of the catalytic subunit of gamma-glutamylcysteine synthetase. This gene encodes a protein containing KELCH-1 like domains, as well as a BTB/POZ domain. Kelch-like ECH-associated protein 1 interacts with NF-E2-related factor 2 in a redox-sensitive manner and the dissociation of the proteins in the cytoplasm is followed by transportation of NF-E2-related factor 2 to the nucleus. This interaction results in the expression of the catalytic subunit of gamma-glutamylcysteine synthetase. Two alternatively spliced transcript variants encoding the same isoform have been found for this gene.

Synonyms: INrf2; KLHL19


Note: Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human:

100%; Mouse: 100%; Bovine: 100%; Rabbit: 100%; Guinea pig: 100%; Zebrafish: 93%

**Protein Families:** Transcription Factors


**Protein Pathways:** Ubiquitin mediated proteolysis

## **Product images:**



WB Suggested Anti-KEAP1 Antibody Titration: 1.0 ug/ml; Positive Control: HepG2 cell lysate KEAP1 is supported by BioGPS gene expression data to be expressed in HepG2






#### Anti-KEAP1 Western Blot & Peptide Block Validation

Lane A: Primary Antibody Lane B: Primary Antibody + Blocking Peptide

Primary Antibody Concentration: 1.25µg/ml Peptide Concentration: 1.0µg/ml Lysate Quantity: 25µg/lane Gel Concentration: 12% Host: Rabbit; Target Name: KEAP1; Sample Tissue: HepG2; Lane A: Primary Antibody; Lane B: Primary Antibody + Blocking Peptide; Primary Antibody Concentration:1.25ug/mL; Peptide Concentration: 1.0ug/mL; Lysate Quantity: 25ug/lane; Gel Concentration: 12%KEAP



Human kidney