

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA337892

NFX1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-NFX1 antibody: synthetic peptide directed towards the N terminal of human NFX1. Synthetic peptide located within the following region: MAEAPPVSGTFKFNTDAAEFIPQEKKNSGLNCGTQRRLDSNRIGRRNYSS
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	113 kDa
Gene Name:	nuclear transcription factor, X-box binding 1
Database Link:	<u>NP_667344</u> <u>Entrez Gene 4799 Human</u> <u>Q12986</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE NFX1 Rabbit Polyclonal Antibody – TA337892

Background:NFX1 binds to the X-box motif of MHC class II genes and represses their expression. NFX1
may play an important role in regulating the duration of an inflammatory response by limiting
the period in which MHC class II molecules are induced by interferon-gamma. Isoform 3
binds to the X-box motif of TERT promoter and represses its expression. Together with
PABPC1 or PABPC4, isoform 1 acts as a coactivator for TERT expression. NFX1 mediates E2-
dependent ubiquitination.MHC class II gene expression is controlled primarily at the
transcriptional level by transcription factors that bind to the X and Y boxes, two highly
conserved elements in the proximal promoter of MHC class II genes. The protein encoded by
this gene is a transcriptional repressor capable of binding to the conserved X box motif of
HLA-DRA and other MHC class II genes in vitro. The protein may play a role in regulating the
duration of an inflammatory response by limiting the period in which class II MHC molecules
are induced by IFN-gamma. Three alternative splice variants, each of which encodes a
different isoform, have been identified.

Synonyms: NFX2; TEG-42; Tex42

 Note:
 Immunogen Sequence Homology: Human: 100%; Dog: 93%; Pig: 93%; Rat: 93%; Horse: 93%;

 Mouse: 93%; Bovine: 93%; Rabbit: 93%; Guinea pig: 85%

Protein Families: Druggable Genome, Transcription Factors

Product images:

168 kDa__ 144 kDa__ 90 kDa__ 65 kDa__ 40 kDa__

WB Suggested Anti-NFX1 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1:62500; Positive Control: Jurkat cell lysate NFX1 is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US