

Product datasheet for **TA338251**

ROR alpha (RORA) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-RORA antibody: synthetic peptide directed towards the middle region of human RORA. Synthetic peptide located within the following region: PGEAELPTPTYNISANGLTELHDDLSNYIDGHTPEGSKADSAVSSFYLDI
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	62 kDa
Gene Name:	RAR related orphan receptor A
Database Link:	NP_002934 Entrez Gene 6095 Human P35398



[View online »](#)

Background:

The protein encoded by this gene is a member of the NR1 subfamily of nuclear hormone receptors. It can bind as a monomer or as a homodimer to hormone response elements upstream of several genes to enhance the expression of those genes. The encoded protein has been shown to interact with NM23-2, a nucleoside diphosphate kinase involved in organogenesis and differentiation, as well as with NM23-1, the product of a tumor metastasis suppressor candidate gene. Also, it has been shown to aid in the transcriptional regulation of some genes involved in circadian rhythm. Four transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2014]

Synonyms:


NR1F1; ROR1; ROR2; ROR3; RZR-ALPHA; RZRA

Note:

Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Goat: 100%; Horse: 100%; Human: 100%; Bovine: 100%; Rabbit: 100%; Rat: 86%; Mouse: 86%; Guinea pig: 86%

Protein Families:

Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Product images:

WB Suggested Anti-RORA Antibody Titration: 0.2-1 ug/ml; Positive Control: HepG2 cell lysate