

Product datasheet for **TA338534**

VDAC2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-VDAC2 antibody: synthetic peptide directed towards the N terminal of human VDAC2. Synthetic peptide located within the following region: MATHGQTCARPMCIPPSYADLGKAARDIFNKGFGFGLVKLDVKTCSGSV
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Protein A purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	32 kDa
Gene Name:	voltage dependent anion channel 2
Database Link:	NP_003366 Entrez Gene 7417 Human P45880



[View online »](#)

Background: This gene encodes a member of the voltage-dependent anion channel pore-forming family of proteins that are considered the main pathway for metabolite diffusion across the mitochondrial outer membrane. The encoded protein is also thought to be involved in the mitochondrial apoptotic pathway via regulation of BCL2-antagonist/killer 1 protein activity. Pseudogenes have been identified on chromosomes 1, 2, 12 and 21, and alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2010]


Synonyms: POR

Note: Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Bovine: 100%; Rabbit: 100%; Guinea pig: 100%; Zebrafish: 86%; Sheep: 79%


Protein Families: Druggable Genome, Ion Channels: Other

Protein Pathways: Calcium signaling pathway, Huntington's disease, Parkinson's disease


Product images:



WB Suggested Anti-VDAC2 Antibody ; Titration: 1.6 ug/ml; Positive Control: human cell lines



WB Suggested Anti-VDAC2 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1:312500; Positive Control: Transfected 293T VDAC2 is supported by BioGPS gene expression data to be expressed in HEK293T



Rabbit Anti-VDAC2 Antibody; Formalin Fixed Paraffin Embedded Tissue: Human Bronchial Epithelial Tissue; Observed Staining: Cytoplasmic; Primary Antibody Concentration: 1:100; Secondary Antibody: Donkey anti-Rabbit-Cy3; Secondary Antibody Concentration: 1