

## Product datasheet for **TA339096**

### PIAS2 Rabbit Polyclonal Antibody


#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-PIAS2 antibody: synthetic peptide directed towards the C terminal of human PIAS2. Synthetic peptide located within the following region: FLDSLTSPLTASSTSVTTTSSHESSTHVSSSSSRSETGVITSSGSNIPDI
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Protein A purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	68 kDa
Gene Name:	protein inhibitor of activated STAT 2
Database Link:	<a href="#">NP_004662</a> <a href="#">Entrez Gene 9063 Human</a> <a href="#">O75928</a>




[View online »](#)

<b>Background:</b>	This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Isoforms of the encoded protein enhance the sumoylation of specific target proteins including the p53 tumor suppressor protein, c-Jun, and the androgen receptor. A pseudogene of this gene is located on the short arm of chromosome 4. The symbol MIZ1 has also been associated with ZBTB17 which is a different gene located on chromosome 1. [provided by RefSeq, Aug 2011]
<b>Synonyms:</b>	ARIP3; DIP; MIZ; MIZ1; PIASX; PIASX-ALPHA; PIASX-BETA; SIZ2; ZMIZ4
<b>Note:</b>	Immunogen Sequence Homology: Dog: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Bovine: 93%
<b>Protein Families:</b>	Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors
<b>Protein Pathways:</b>	Jak-STAT signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated proteolysis

**Product images:**


WB Suggested Anti-PIAS2 Antibody Titration: 0.6 ug/ml; ELISA Titer: 1:312500; Positive Control: HepG2 cell lysate

## PIAS2




Host: Rabbit; Target Name: PIAS2; Sample Tissue: HeLa; Antibody Dilution: 1.0 ug/ml

## PIAS2




Host: Rabbit; Target Name: PIAS2; Sample Tissue: MCF7; Antibody Dilution: 1.0 ug/ml

## PIAS2



Host: Rabbit; Target Name: PIAS2; Sample Tissue: HepG2; Antibody Dilution: 1.0 ug/ml

## PIAS2



Rabbit Anti-PIAS2  
Sample Type: 721\_B  
Antibody Concentration: 1ug/mL

Host: Rabbit; Target Name: PIAS2; Sample Tissue: 721\_B; Antibody Dilution: 1.0 ug/ml PIAS2 is strongly supported by BioGPS gene expression data to be expressed in Human 721\_B cells