

Product datasheet for **TA339097**

N myc interactor (NMI) Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-NMI antibody: synthetic peptide directed towards the N terminal of human NMI. Synthetic peptide located within the following region: ETKMKFLSVETPENDSQLSNISCSFQVSSKVPYEIQKGQALITFEKEEVA
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Protein A purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	35 kDa
Gene Name:	N-myc and STAT interactor
Database Link:	NP_004679 Entrez Gene 9111 Human Q13287



[View online »](#)

- Background:** NMYC interactor (NMI) encodes a protein that interacts with NMYC and CMYC (two members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. The NMI protein also interacts with all STATs except STAT2 and augments STAT-mediated transcription in response to cytokines IL2 and IFN-gamma. The NMI mRNA has low expression levels in all human fetal and adult tissues tested except brain and has high expression in cancer cell line-myeloid leukemias. [provided by RefSeq, Jul 2008]
- Synonyms:** N-myc (and STAT) interactor; N-myc-interactor; N-myc and STAT interactor; N-myc interactor
- Note:** Immunogen Sequence Homology: Human: 100%; Pig: 93%; Bovine: 93%; Dog: 90%; Horse: 90%; Mouse: 90%; Guinea pig: 90%; Rabbit: 83%; Rat: 77%
- Protein Families:** Druggable Genome, Transcription Factors

Product images:

WB Suggested Anti-NMI Antibody Titration: 1.25 ug/ml; ELISA Titer: 1:62500; Positive Control: Human Liver