

## **Product datasheet for TA339209**

## **OAS1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications:IHC, WBRecommended Dilution:WB, IHCReactivity:HumanHost:RabbitIsotype:IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-OAS1 antibody: synthetic peptide directed towards the middle

region of human OAS1. Synthetic peptide located within the following region:

RRQLTKPRPVILDPADPTGNLGGGDPKGWRQLAQEAEAWLNYPCFKNWDG

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

Purification: Protein A purified

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 47 kDa

**Gene Name:** 2'-5'-oligoadenylate synthetase 1

Database Link: NP 001027581

Entrez Gene 4938 Human

P00973



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

This protein is a member of the 2-5A synthetase family, essential proteins involved in the innate immune response to viral infection. The encoded protein is induced by interferons and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize 2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral RNA degradation and the inhibition of viral replication. This gene encodes a member of the 2-5A synthetase family, essential proteins involved in the innate immune response to viral infection. The encoded protein is induced by interferons and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize 2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral RNA degradation and the inhibition of viral replication. The three known members of this gene family are located in a cluster on chromosome 12. Mutations in this gene have been associated with host susceptibility to viral infection. Alternatively spliced transcript variants encoding different isoforms have been described.


Synonyms: IFI-4; OIAS; OIASI

Note: Immunogen Sequence Homology: Human: 100%; Dog: 93%; Pig: 93%; Rat: 93%; Horse: 93%;

Mouse: 93%; Bovine: 93%; Guinea pig: 87%; Rabbit: 86%


**Protein Families:** Druggable Genome

## **Product images:**



WB Suggested Anti-OAS1 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 312500; Positive Control: Transfected 293T





Immunohistochemistry with Small intestine tissue