

## **Product datasheet for TA339586**

## **PCDHA6 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** The immunogen for anti-PCDHA6 antibody: synthetic peptide directed towards the C terminal

of human PCDHA6. Synthetic peptide located within the following region:

LVKDHGEPALTATATVLVSLVESGQAPKASSRASVGAAGPEAALVDVNVY

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

**Purification:** Protein A purified

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 84 kDa

**Gene Name:** protocadherin alpha 6

Database Link: NP 114036

Entrez Gene 56142 Human

Q9UN73



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

PCDHA6 contains 6 cadherin domains. It is a potential calcium-dependent cell-adhesion protein. PCDHA6 may be involved in the establishment and maintenance of specific neuronal connections in the brain. This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined.


Synonyms: CNR2; CNRN2; CNRS2; CRNR2; PCDH-ALPHA6

Note: Immunogen Sequence Homology: Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Guinea

pig: 100%; Mouse: 93%; Dog: 86%; Bovine: 86%

**Protein Families:** Secreted Protein, Transmembrane

## **Product images:**



WB Suggested Anti-PCDHA6 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 62500; Positive

Control: 293T cell lysate