

Product datasheet for **TA340190**

gamma C Crystallin (CRYGC) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-CRYGC antibody: synthetic peptide directed towards the middle region of human CRYGC. Synthetic peptide located within the following region: GLSDSIRSCCLIPQTVSHRLRLYEREDHKGLMMELSEDCPSIQDRFHLSE
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	21 kDa
Gene Name:	crystallin gamma C
Database Link:	NP_066269 Entrez Gene 1420 Human P07315



[View online »](#)

Background:

Crystallins are the dominant structural components of the vertebrate eye lens. Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. Four gamma-crystallin genes (gamma-A through gamma-D) and three pseudogenes (gamma-E, gamma-F, gamma-G) are tandemly organized in a genomic segment as a gene cluster. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Synonyms:

CCL; CRYG3; CTRCT2

Note:

Immunogen Sequence Homology: Human: 100%; Dog: 93%; Horse: 86%; Rabbit: 86%; Rat: 79%; Bovine: 79%

Protein Families:

Druggable Genome

Product images:

WB Suggested Anti-CRYGC Antibody Titration:
0.2-1 ug/ml; ELISA Titer: 1: 62500; Positive
Control: OVCAR-3 cell lysate
CRYGC is supported
by BioGPS gene expression data to be expressed
in OVCAR3