

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA341926

### **PCDHA4 Rabbit Polyclonal Antibody**

### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-PCDHA4 antibody: synthetic peptide directed towards the C terminal of human PCDHA4. Synthetic peptide located within the following region: SGYNAWLSYELQPETASASIPFRVGLYTGEISTTRALDETDAPRQRLLVL
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	99 kDa
Gene Name:	protocadherin alpha 4
Database Link:	<u>NP_061730</u> <u>Entrez Gene 56144 Human</u> <u>Q9UN74</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **PCDHA4** Rabbit Polyclonal Antibody – TA341926


Background:	This gene is a member ofThe protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related toThe mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes inThe cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains whileThe C-terminal exons encodeThe cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role inThe establishment and function of specific cell-cell connections inThe brain. Alternative splicing has been observed and additional variants have been suggested butTheir full-length nature has yet to be determined. [provided by RefSeq, Jul 2008]
Synonyms:	CNR1; CNRN1; CRNR1; PCDH-ALPHA4
NISter	L

 Note:
 Immunogen Sequence Homology: Pig: 100%; Human: 100%; Guinea pig: 100%; Dog: 93%; Rat:

 93%; Horse: 93%; Mouse: 93%; Bovine: 93%

Protein Families: Transmembrane

# **Product images:**



WB Suggested Anti-PCDHA4 Antibody Titration: 0.2-1 ug/ml; Positive Control: Hela cell lysate

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US