

Product datasheet for TA342550

NDRG1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-NDRG1 antibody: synthetic peptide directed towards the N terminal

of human NDRG1. Synthetic peptide located within the following region: MSREMQDVDLAEVKPLVEKGETITGLLQEFDVQEQDIETLHGSVHVTLCG

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43 kDa

Gene Name: N-myc downstream regulated 1

Database Link: NP 006087

Entrez Gene 10397 Human

Q92597

Background: This gene is a member of the N-myc downregulated gene family which belongs to the

alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein

involved in stress responses, hormone responses, cell growth, and differentiation. The encoded protein is necessary for p53-mediated caspase activation and apoptosis. Mutations

in this gene are a cause of Charcot-Marie-Tooth disease type 4D, and expression of this gene may be a prognostic indicator for several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq,

May 2012]



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: CAP43; CMT4D; DRG-1; DRG1; GC4; HMSNL; NDR1; NMSL; PROXY1; RIT42; RTP; TARG1; TDD5

Note: Immunogen Sequence Homology: Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Mouse:

100%; Bovine: 100%; Rabbit: 100%; Dog: 93%; Guinea pig: 93%

Product images:



WB Suggested Anti-NDRG1 Antibody Titration: 2.5 ug/ml; Positive Control: HepG2 cell lysate