

Product datasheet for TA344764

WDR35 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-WDR35 antibody: synthetic peptide directed towards the N terminal

of human WDR35. Synthetic peptide located within the following region: SGSVQVVTWNEQYQKLTTSDENGLIIVWMLYKGSWIEEMINNRNKSVVRS

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Affinity Purified

Conjugation: Unconjugated

Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 132 kDa

Gene Name: WD repeat domain 35

Database Link: NP 065830

Entrez Gene 57539 Human

Q9P2L0



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



WDR35 Rabbit Polyclonal Antibody - TA344764

Background: This gene encodes a member of the WD repeat protein family. WD repeats are minimally

conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene, but the


biological validity of some variants has not been determined.

Synonyms: CED2; IFT121; IFTA1; SRTD7

Note: Immunogen Sequence Homology: Human: 100%; Dog: 93%; Pig: 93%; Rat: 93%; Horse: 93%;

Mouse: 93%; Rabbit: 93%; Zebrafish: 93%; Guinea pig: 93%

Product images:



WB Suggested Anti-WDR35 Antibody Titration: 0.2-1 ug/ml; Positive Control: Human Lung