

Product datasheet for **TA345815**

RBMS1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB, IHC
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-RBMS1 antibody: synthetic peptide directed towards the C terminal of human RBMS1. Synthetic peptide located within the following region: TYMPATSAMQGAYLPQYAHMQTTAVPVEEASGQQQVAVETSNDHSPYTFQ
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Purification:	Protein A purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	44 kDa
Gene Name:	RNA binding motif single stranded interacting protein 1
Database Link:	NP_002888 Entrez Gene 5937 Human P29558



[View online »](#)

Background:


RBMS1 is a member of a small family of proteins which bind single stranded DNA/RNA. These proteins are characterized by the presence of two sets of ribonucleoprotein consensus sequence (RNP-CS) that contain conserved motifs, RNP1 and RNP2, originally described in RNA binding proteins, and required for DNA binding. These proteins have been implicated in such diverse functions as DNA replication, gene transcription, cell cycle progression and apoptosis. This gene encodes a member of a small family of proteins which bind single stranded DNA/RNA. These proteins are characterized by the presence of two sets of ribonucleoprotein consensus sequence (RNP-CS) that contain conserved motifs, RNP1 and RNP2, originally described in RNA binding proteins, and required for DNA binding. These proteins have been implicated in such diverse functions as DNA replication, gene transcription, cell cycle progression and apoptosis. Multiple transcript variants, resulting from alternative splicing and encoding different isoforms, have been described. Several of these were isolated by virtue of their binding to either strand of an upstream element of c-myc (MSSPs), or by phenotypic complementation of cdc2 and cdc13 mutants of yeast (scr2), or as a potential human repressor of HIV-1 and ILR-2 alpha promoter transcription (YC1). A pseudogene for this locus is found on chromosome 12.

Synonyms:


C2orf12; HCC-4; MSSP; MSSP-1; MSSP-2; MSSP-3; SCR2; YC1

Note:

Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Guinea pig: 100%; Goat: 93%; Bovine: 93%

Product images:

WB Suggested Anti-RBMS1 Antibody Titration: 2.5 ug/ml; Positive Control: Jurkat cell lysate



Rabbit Anti-RBMS1 Antibody; Paraffin Embedded Tissue: Human Skin; Cellular Data: Squamous epithelial cells; Antibody Concentration: 4.0-8.0 ug/ml; Magnification: 400X