

Product datasheet for **TA345905**

PCBP2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB, IHC
Reactivity:	Mouse, Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-PCBP2 antibody: synthetic peptide directed towards the middle region of human PCBP2. Synthetic peptide located within the following region: VIFAGGQDRYSTGSDSASFHTTPSMCLNPDLEGPPLEAYTIQQQYAI PQ
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Purification:	Protein A purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	40 kDa
Gene Name:	poly(rC) binding protein 2
Database Link:	NP_005007 Entrez Gene 18521 Mouse Entrez Gene 5094 Human Q15366



[View online »](#)

Background:


PCBP2 appears to be multifunctional. It along with PCBP-1 and hnRNPk corresponds to the major cellular poly(rC)-binding proteins. This protein together with PCBP-1 also functions as translational coactivators of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES and promote poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human Papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This gene and PCBP-1 has paralogues PCBP3 and PCBP4 which is thought to arose as a result of duplication events of entire genes. The protein encoded by this gene appears to be multifunctional. It along with PCBP-1 and hnRNPk corresponds to the major cellular poly(rC)-binding proteins. It contains three K-homologous (KH) domains which may be involved in RNA binding. This encoded protein together with PCBP-1 also functions as translational coactivators of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES and promote poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human Papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1 intronless gene which has similar functions. This gene and PCBP-1 has paralogues PCBP3 and PCBP4 which is thought to arose as a result of duplication events of entire genes. It also has two processed pseudogenes PCBP2P1 and PCBP2P2. There are presently two alternatively spliced transcript variants described for this gene.

Synonyms:


hnRNP-E2; HNRNPE2; HNRPE2

Note:

Immunogen Sequence Homology: Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Bovine: 100%; Guinea pig: 100%

Product images:

WB Suggested Anti-PCBP2 Antibody Titration: 1.25 ug/ml; Positive Control: Jurkat cell lysate There is BioGPS gene expression data showing that PCBP2 is expressed in Jurkat



Rabbit Anti-PCBP2 Antibody; Paraffin Embedded Tissue: Human Lung; Cellular Data: Alveolar cells; Antibody Concentration: 4.0-8.0 ug/ml; Magnification: 400X