

## **Product datasheet for TA346151**

## **ANK1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

**Isotype:** IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-ANK1 antibody: synthetic peptide directed towards the middle

region of human ANK1. Synthetic peptide located within the following region:

PCAMPETVVIRSEEQEQASKEYDEDSLIPSSPATETSDNISPVASPVHTG

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Purification:** Affinity Purified

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 189 kDa
Gene Name: ankyrin 1

Database Link: NP 065210

Entrez Gene 286 Human

P16157



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Ankyrins are a family of proteins that link the integral membrane proteins to the underlying spectrin-actin cytoskeleton and play key roles in activities such as cell motility, activation, proliferation, contact and the maintenance of specialized membrane domains. Multiple isoforms of ankyrin with different affinities for various target proteins are expressed in a tissue-specific, developmentally regulated manner. Most ankyrins are typically composed of three structural domains: an amino-terminal domain containing multiple ankyrin repeats; a central region with a highly conserved spectrin binding domain; and a carboxy-terminal regulatory domain which is the least conserved and subject to variation. Ankyrin 1, the prototype of this family, was first discovered in the erythrocytes, but since has also been found in brain and muscles. Mutations in erythrocytic ankyrin 1 have been associated in approximately half of all patients with hereditary spherocytosis. Complex patterns of alternative splicing in the regulatory domain, giving rise to different isoforms of ankyrin 1 have been described. Truncated muscle-specific isoforms of ankyrin 1 resulting from usage of an alternate promoter have also been identified. [provided by RefSeq, Dec 2008]


**Synonyms:** ANK; SPH1; SPH2

Note: Immunogen Sequence Homology: Human: 100%; Dog: 93%; Pig: 93%; Rat: 93%; Mouse: 93%;

Rabbit: 93%; Guinea pig: 93%; Horse: 92%

**Protein Families:** Transmembrane


## **Product images:**



WB Suggested Anti-ANK1 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 12500; Positive Control: HT1080 cell lysate







Host: Rabbit; Target Name: ANK1; Sample Tissue: Human Fetal Liver; Antibody Dilution: 1.0 ug/ml