

Product datasheet for TA346193

ABCB4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-ABCB4 antibody: synthetic peptide directed towards the N terminal

of human ABCB4. Synthetic peptide located within the following region:

AGAVAEEALGAIRTVIAFGGQNKELERYQKHLENAKEIGIKKAISANISM

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 141 kDa

Gene Name: ATP binding cassette subfamily B member 4

Database Link: NP 000434

Entrez Gene 5244 Human

P21439



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

ABCB4, a membrane-associated protein, is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance as well as antigen presentation. ABCB4 is a full transporter and member of the p-glycoprotein family of membrane proteins with phosphatidylcholine as its substrate. The function of this protein has not yet been determined; however, it may involve transport of phospholipids from liver hepatocytes into bile. The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance as well as antigen presentation. This gene encodes a full transporter and member of the p-glycoprotein family of membrane proteins with phosphatidylcholine as its substrate. The function of this protein has not yet been determined; however, it may involve transport of phospholipids from liver hepatocytes into bile. Alternative splicing of this gene results in several products of undetermined function.

Synonyms: 3; ABC21; GBD1; ICP3; MDR2; MDR3; PFIC-3; PGY3


Note: Immunogen Sequence Homology: Rat: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Pig:

93%; Bovine: 86%; Dog: 77%; Horse: 77%; Sheep: 77%; Guinea pig: 77%

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: ABC transporters

Product images:



WB Suggested Anti-ABCB4 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 312500; Positive Control: 293T cell lysate





Anti-ABCB4 Western Blot & Peptide Block Validation

Lysate: 293T Cell

Lane A: Primary Antibody Lane B: Primary Antibody + Blocking Peptide

Primary Antibody Concentration: 1.0µg/ml Peptide Concentration: 5.0µg/ml Lysate Quantity: 25µg/lane Gel Concentration: 12% Host: Rabbit; Target Name: ABCB4; Sample Tissue: 293T Whole Cell; Lane A: Primary Antibody; Lane B: Primary Antibody + Blocking Peptide; Primary Antibody Concentration: 1 ug/ml; Peptide Concentration: 5 ug/ml; Lysate Quantity: 25 ug/lane/Lane; Gel Concentration: 0.12