

Product datasheet for TA346220

CYP3A7 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-CYP3A7 antibody: synthetic peptide directed towards the middle

region of human CYP3A7. Synthetic peptide located within the following region:

KLALVRVLQNFSFKPCKETQIPLKLRFGGLLLTEKPIVLKAESRDETVSG

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Affinity Purified

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 57 kDa

Gene Name: cytochrome P450 family 3 subfamily A member 7

Database Link: NP 000756

Entrez Gene 1551 Human

P24462



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

CYP3A7 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This enzyme hydroxylates testosterone and dehydroepiandrosterone 3-sulphate, which is involved in the formation of estriol during pregnancy. The enzyme also metabolizes some drugs such as aflatoxin B1. This gene, CYP3A7, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This enzyme hydroxylates testosterone and dehydroepiandrosterone 3-sulphate, which is involved in the formation of estriol during pregnancy. The enzyme also metabolizes some drugs such as aflatoxin B1. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Transcript variants have been described, but it is not known whether these transcripts are normally produced.

Synonyms: CP37; CYPIIIA7; P-450(HFL33); P-450111A7; P450-HFLA

Note: Immunogen Sequence Homology: Human: 100%; Horse: 86%


Protein Families: Druggable Genome, P450, Transmembrane

Protein Pathways: Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Linoleic acid

metabolism, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol

metabolism

Product images:



WB Suggested Anti-CYP3A7 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 1562500; Positive

Control: HT1080 cell lysate