

## **Product datasheet for TA346445**

# **ZNF19 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** The immunogen for anti-ZNF19 antibody: synthetic peptide directed towards the N terminal

of human ZNF19. Synthetic peptide located within the following region: TALGYPVPKPALISLLERGDMAWGLEAQDDPPAERTKNVCKDVETNIDSE

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Purification:** Affinity Purified

**Conjugation:** Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 52 kDa

**Gene Name:** zinc finger protein 19

Database Link: NP 008892

Entrez Gene 7567 Human

P17023



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **ZNF19 Rabbit Polyclonal Antibody - TA346445**

**Background:** ZNF19 contains a zinc finger, a nucleic acid-binding domain present in many transcription

factors. The protein encoded by this gene contains a zinc finger, a nucleic acid-binding domain present in many transcription factors. This gene is located in a region next to ZNF23, a gene also encoding a zinc finger protein, on chromosome 16. The protein encoded by this gene contains a zinc finger, a nucleic acid-binding domain present in many transcription factors. This gene is located in a region next to ZNF23, a gene also encoding a zinc finger


protein, on chromosome 16.

Synonyms: KOX12

**Note:** Immunogen Sequence Homology: Human: 100%; Horse: 79%

**Protein Families:** Transcription Factors

### **Product images:**



WB Suggested Anti-ZNF19 Antibody Titration: 0.2-1 ug/ml; Positive Control: Jurkat cell lysate