

## **Product datasheet for TA346475**

## **FADS1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** The immunogen for anti-FADS1 antibody: synthetic peptide directed towards the N terminal

of human FADS1. Synthetic peptide located within the following region: RHPGGSRVISHYAGQDATDPFVAFHINKGLVKKYMNSLLIGELSPEQPSF

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Purification:** Affinity Purified

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 52 kDa

**Gene Name:** fatty acid desaturase 1

Database Link: NP 037534

Entrez Gene 3992 Human

O60427



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

FADS1 is a member of the fatty acid desaturase (FADS) family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, both of which are characterized by conserved histidine motifs. The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, both of which are characterized by conserved histidine motifs. This gene is clustered with family members FADS1 and FADS2 at 11q12-q13.1; this cluster is thought to have arisen evolutionarily from gene duplication based on its similar exon/intron organization. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.


**Synonyms:** D5D; FADS6; FADSD5; LLCDL1; TU12

**Note:** Immunogen Sequence Homology: Pig: 100%; Rat: 100%; Human: 100%; Mouse: 100%; Bovine:

100%; Dog: 93%; Horse: 83%; Rabbit: 83%; Guinea pig: 83%

Protein Families: Transcription Factors, Transmembrane
Protein Pathways: Biosynthesis of unsaturated fatty acids

## **Product images:**



WB Suggested Anti-FADS1 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 312500; Positive Control: 721\_B cell lysate; FADS1 is strongly supported by BioGPS gene expression data to be expressed in Human 721 B cells