

Product datasheet for TA346564

Decorin (DCN) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:IHC, WBRecommended Dilution:WB, IHCReactivity:HumanHost:RabbitIsotype:IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-DCN antibody: synthetic peptide directed towards the N terminal of

human DCN. Synthetic peptide located within the following region: IGPEVPDDRDFEPSLGPVCPFRCQCHLRVVQCSDLGLDKVPKDLPPDTTL

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Affinity Purified
Conjugation: Unconjugated

Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36 kDa

Gene Name: decorin

Database Link: NP 001911

Entrez Gene 1634 Human

P07585



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: The protein encoded by this gene is a small cellular or pericellular matrix proteoglycan that is

closely related in structure to biglycan protein. The encoded protein and biglycan are thought to be the result of a gene duplication. This protein is a component of connective tissue, binds

to type I collagen fibrils, and plays a role in matrix assembly. It contains one attached

glycosaminoglycan chain. This protein is capable of suppressing the growth of various tumor cell lines. There are multiple alternatively spliced transcript variants known for this gene. This

gene is a candidate gene for Marfan syndrome. [provided by RefSeq, Jul 2008]

Synonyms: CSCD; DSPG2; PG40; PGII; PGS2; SLRR1B


Note: Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Horse: 100%; Human: 100%; Rabbit:

100%; Mouse: 93%; Goat: 92%; Sheep: 92%; Bovine: 92%; Rat: 86%; Guinea pig: 86%


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: TGF-beta signaling pathway

Product images:



WB Suggested Anti-DCN Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 312500; Positive Control: Human Placenta



Skin