

## Product datasheet for **TA346674**

### GSTA3 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-GSTA3 antibody: synthetic peptide directed towards the middle region of human GSTA3. Synthetic peptide located within the following region: SSLISNFPLLKALKTRISNLPTVKKFLQPGSPRKPPADAKALEEARKIFR
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25 kDa
Gene Name:	glutathione S-transferase alpha 3
Database Link:	<a href="#">NP_000838</a> <a href="#">Entrez Gene 2940 Human</a> <a href="#">Q16772</a>



[View online »](#)

**Background:**

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes are involved in cellular defense against toxic, carcinogenic, and pharmacologically active electrophilic compounds. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase belonging to the alpha class genes that are located in a cluster mapped to chromosome 6. Genes of the alpha class are highly related and encode enzymes with glutathione peroxidase activity. However, during evolution, this alpha class gene diverged accumulating mutations in the active site that resulted in differences in substrate specificity and catalytic activity. The enzyme encoded by this gene catalyzes the double bond isomerization of precursors for progesterone and testosterone during the biosynthesis of steroid hormones. An additional transcript variant has been identified, but its full length sequence has not been determined. [provided by RefSeq, Jul 2008]

**Synonyms:**


GSTA3-3; GTA3

**Note:**

Immunogen Sequence Homology: Human: 100%; Mouse: 79%; Sheep: 77%

**Protein Pathways:**

Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by cytochrome P450

**Product images:**


WB Suggested Anti-GSTA3 Antibody Titration: 0.2-1 ug/ml; Positive Control: HCT15 cell lysate