

**Name: Mouse Monoclonal AKT(pan) Antibody**  
**Product Data Sheet - ANTIBODY****Catalog: TA346875**

<b>Components:</b>	• Mouse Monoclonal AKT(pan) Antibody (TA346875)
<b>Amount:</b>	100ul
<b>Immunogen:</b>	The immunogen for AKT antibody: purified recombinant human AKT1 protein fragments expressed in E.coli.
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG2b
<b>Species Reactivity:</b>	M,R
<b>Guaranteed Applications:</b>	WB
<b>Suggested Dilutions:</b>	WB: 1:1000
<b>Concentration:</b>	Lot dependent; please refer to CoA along with shipment
<b>Buffer:</b>	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.02% sodium azide, 50%,glycerol
<b>Purification:</b>	Affinity purified
<b>Storage Condition:</b>	Shipped at -20C or with ice packs. Upon delivery store at -20C. Dilute in PBS (pH7.3) if necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

## Target

<b>Target Name:</b>	AKT serine/threonine kinase 1
<b>Alternative Name:</b>	AKT CWS6 PKB PKB-ALPHA PRKBA RAC RAC-ALPHA
<b>Database Link:</b>	<a href="#">NP_001014432</a>
<b>Function:</b>	Akt1 is involved in cellular survival pathways,by inhibiting apoptotic processes.Akt1 is also able to induce protein synthesis pathways,and is therefore a key signaling protein in the cellular pathways that lead to skeletal muscle hypertrophy, and general tissue growth.Since it can block apoptosis,and thereby promote cell survival,Akt1 has been implicated as a major factor in many types of cancer.Akt(now also called Akt1)was originally identified as the oncogene in the transforming retrovirus,AKT8. Akt2 is an important signaling molecule in the Insulin signaling pathway.It is required to induce glucose transport.In a mouse which is for Akt1 but normal for Akt2,glucose homeostasis is unperturbed,but the animals are

***This product is to be used for laboratory only. Not for diagnostic or therapeutic use.***

smaller, consistent with a role for Akt1 in growth. In contrast, mice which do not have Akt2, but have normal Akt1, have mild growth deficiency and display a diabetic phenotype (insulin resistance), again consistent with the idea that Akt2 is more specific for the insulin receptor signaling pathway. The role of Akt3 is less clear, though it appears to be predominantly expressed in the brain. It has been reported that mice lacking Akt3 have small brains.


---

***This product is to be used for laboratory only. Not for diagnostic or therapeutic use.***


© 2017 OriGene Technologies, Inc. 9620 Medical Center Dr., Suite 200, Rockville, MD 20850

---

## Validation Data



Western blot detection of total AKT in rat heart, rat testis, mouse brain, mouse spleen and C6 cell lysates and using AKT(pan) mouse mAb (1:1000 diluted). Predicted band size: 60KDa. Observed band size: 60KDa.



L: Western blot detection of DSBA in AKT1, AKT2, AKT3 and DSBA recombinant antigen fragments the same sample quality, and using DSBA mouse mAb (1:1000 diluted). R: Western blot detection of AKT in AKT1, AKT2 and AKT3 recombinant antigen fragments and using AKT(pan) mouse mAb (1:1000 diluted).

\* More validation images may be available on our website: <http://www.origene.com/antibody/TA346875.aspx>

***This product is to be used for laboratory only. Not for diagnostic or therapeutic use.***

© 2017 OriGene Technologies, Inc. 9620 Medical Center Dr., Suite 200, Rockville, MD 20850