

Product datasheet for TA346883M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

AMACR Mouse Monoclonal Antibody [Clone ID: 2F3-C11-B12]

Product data:

Product Type: Primary Antibodies

Clone Name: 2F3-C11-B12

Applications: IF, WB

Recommended Dilution: WB: 1:1000, IF: 1:100

Reactivity: Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: The immunogen for AMACR antibody: purified recombinant human AMACR(C-terminus)

protein fragments expressed in E.coli.

Formulation: Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with

0.02% sodium azide, 50%, glycerol

Purification: Affinity purified
Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42 kDa

Gene Name: alpha-methylacyl-CoA racemase

Database Link: NP 055139

Entrez Gene 25284 Rat

Q9UHK6

Background: This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-

bile acylCoAs between their (R)-and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy,

pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis.

Alternatively spliced transcript variants have been described.






Synonyms: AMACRD; CBAS4; RACE; RM


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Primary bile acid biosynthesis

Product images:



Western blot detection of AMACR (C-terminus) in Rat Liver lysates using AMACR (C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 42KDa. Observed band size: 42KDa.



Immunocytochemistry staining of Jurkat cells fixed with -20? Ethanol and using anti-AMACR (Cterminus) mouse mAb (dilution 1:100).