

Product datasheet for TA346935

E2F1 Mouse Monoclonal Antibody [Clone ID: 4G8-2C2-C12]

Product data:

Product Type:	Primary Antibodies
Clone Name:	4G8-2C2-C12
Applications:	IF, IP, WB
Recommended Dilution:	WB: 1:500, IF: 1:100
Reactivity:	Human, Rat
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	The immunogen for E2F1 antibody: purified recombinant human E2F-1 protein fragments expressed in E.coli.
Formulation:	Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.03% Proclin300 and 50% glycerol.
Purification:	Affinity purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	E2F transcription factor 1
Database Link:	<u>NP_005216</u> <u>Entrez Gene 1869 Human</u> <u>Q01094</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	E2F1 Mouse Monoclonal Antibody [Clone ID: 4G8-2C2-C12] – TA346935
Background:	The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis.
Synonyms:	E2F-1; RBAP1; RBBP3; RBP3
Protein Families	: Druggable Genome, Transcription Factors
Protein Pathwa	ys: Bladder cancer, Cell cycle, Chronic myeloid leukemia, Glioma, Melanoma, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US