

Product datasheet for TA347114

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DNMT3A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: ELISA (1:50); Western blotting (1:2,000); Immunoprecipitation (2 μg/IP)

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-DNMT3A antibody: human DNMT3A, (DNA methyltransferase 3A),

using a KLH-conjugated synthetic peptide corresponding to amino acids 107-121.

Concentration: lot specific

Purification: Affinity purified polyclonal antibody in PBS containing 0.05% azide and 0.05% ProClin 300.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: DNA (cytosine-5-)-methyltransferase 3 alpha

Database Link: NP 783328

Entrez Gene 1788 Human

Q9Y6K1

Background: DNMT3A (UniProtKB/Swiss-Prot entry Q9Y6K1) catalyses the genome wide de novo

methylation of CpG residues. DNA methylation on CpG residues by DNMT3A regulates gene expression and is essential for development. DNMT3A is strongly expressed in embryonic stem cells, but low in adult somatic cells. DNA methylation is coordinated with methylation of

histones. DNMT3A binds to SETDB1 and HDAC1, and is involved in the repression of

transcription from promoters containing an E2F binding site.

Synonyms: DNMT3A2; M.HsallIA; TBRS


Protein Families: Druggable Genome

Protein Pathways: Cysteine and methionine metabolism, Metabolic pathways






Product images:



WB analysis with anti-GAL4 antibody (lane 1 and 2 respectively). Alternatively, GAL4-DNMT3A was immunoprecipitated and WB analysis was performed with the DNMT3A antibody (diluted 1:2,000) (lane 3).



Determination of the antibody titer To determine the titer of the antibody, an ELISA was performed using a serial dilution of the antibody against human DNMT3A, crude serum and Flow Through in antigen coated wells. By plotting the absorbance against the antibody dilution (Figure 1), the titer of the antibody was estimated to be 1:400.