

Product datasheet for **TA347217**

H4-16 Rabbit Polyclonal Antibody

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | Dot, ELISA, IF, WB |
| Recommended Dilution: | ChIP (1ug/IP); ELISA (1:100); Dot blotting (1:20,000); Western blotting (1:200); Immunofluorescence (1:500) |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | The immunogen for anti-H4K8ac antibody: the region of histone H4 containing the acetylated lysine 8 (H4K8ac), using a KLH-conjugated synthetic peptide. |
| Concentration: | lot specific |
| Purification: | Affinity purified polyclonal antibody in PBS containing 0.05% azide and 0.05% ProClin 300. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | histone cluster 4, H4 |
| Database Link: | NP_778224 Entrez Gene 320332 Mouse Entrez Gene 121504 Human P62805 |
| Background: | Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases. |




[View online »](#)


Synonyms: H4; p

Protein Pathways: Systemic lupus erythematosus


Product images:




WB using the antibody against H4K8ac diluted 1:200 in TBS-Tween containing 5% skimmed milk. The position of the protein of interest is indicated on the right; the marker (in kDa) is shown on the left.



Determination of the antibody titer To determine the titer of the antibody, an ELISA was performed using a serial dilution of the antibody against H4K8ac, crude serum and flow through in antigen coated wells. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution (Figure 2), the titer of the purified antibody was estimated to be 1:16,700.



INIH3T3 cells were stained with the ab against H4K8ac and with DAPI. Cells were fixed with 4% formaldehyde for 10' and blocked with PBS/TX-100 containing 5% normal goat serum and 1% BSA. The cells were immunofluorescently labelled with the H4K8ac antibody (left) diluted 1:500 in blocking solution followed by an anti-rabbit antibody conjugated to Alexa488. The middle panel shows staining of the nuclei with DAPI. A merge of the two stainings is shown on the right.



A Dot Blot analysis was performed to test the cross reactivity of the antibody against H4K8ac with peptides containing other histone modifications and the unmodified H4K8. One hundred to 0.2 pmol of the respective peptides were spotted on a membrane. The antibody was used at a dilution of 1:20,000. Image shows a high specificity of the antibody for the modification of interest.