

Product datasheet for TA347290

Froduct datasireet for TAS47230

SIRT1 Mouse Monoclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: Western blotting (1:1,000)

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: The immunogen for anti-SirT1 antibody: human SirT1 (NAD-dependent deacetylase sirtuin-1),

using a synthetic peptide.

Concentration: lot specific

Purification: Ascites fluid from mouse containing 0.05% azide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: sirtuin 1

Database Link: NP 036370

Entrez Gene 23411 Human

Q96EB6

Background: SirT1 (UniProtKB/Swiss-Prot entry Q96EB6) is a NAD-dependent deacetylase, which regulates

processes such as apoptosis and muscle differentiation by the deacetylation of key regulatory proteins. Deacetylation of lysine 382 of p53/TP53 by Sirt1 impairs its ability to induce the proapoptotic program and modulate cell senescence, whereas deacetylation of TAF1B represses rDNA transcription by the RNA polymerase I. SirT1 is also involved in HES1- and HEY2-mediated transcriptional repression and inhibits skeletal muscle differentiation by deacetylating PCAF and MYOD1. SirT1 is recruited to the nuclear bodies via its interaction

with PML.

Synonyms: SIR2; SIR2alpha; SIR2L1

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:

Figure 1



WB was performed using nuclear extracts from HeLa cells (HeLa NE, 20 ug) and the antibody against hSirT1 diluted 1:1,000 in TBS-Tween containing 5% skimmed milk. The molecular weight marker (in kDa) is shown on the left, the position of the protein of interest is shown on the right.