

Product datasheet for TA348928

TIM 1 (HAVCR1) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 0.3-1 ug/ml

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for Anti-IL17C (aa160-172) Antibody: Peptide with sequence

RRPCSRDGSGLPT, from the internal region of the protein sequence according to

NP_037410.1.

Formulation: Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -

20°C. Minimize freezing and thawing.

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: hepatitis A virus cellular receptor 1

Database Link: NP 036338

Entrez Gene 26762 Human

Q96D42

Background: The protein encoded by this gene is a T cell-derived cytokine that shares the sequence

similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine

was found to be restricted to activated T cells. [provided by RefSeq, Jul 2008]

Synonyms: HAVCR; HAVCR-1; KIM-1; KIM1; TIM; TIM-1; TIMD-1; TIMD1

Protein Families: Druggable Genome, Transmembrane




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



TA348928 (0.3 ug/ml) staining of Human Lymph Node lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.