

Product datasheet for **TA348938**

AMHR2 Goat Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1-3 ug/ml
Reactivity:	Human
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for Anti-AMHR2 (aa88-99) Antibody: Peptide with sequence ESLHCDPSRAH, from the internal region of the protein sequence according to NP_065434.1; NP_001158162.1; NP_001158163.1.
Formulation:	Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Concentration:	lot specific
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	anti-Mullerian hormone receptor type 2
Database Link:	NP_065434 Entrez Gene 269 Human Q16671



[View online »](#)

- Background:** This gene encodes the receptor for the anti-Mullerian hormone (AMH) which, in addition to testosterone, results in male sex differentiation. AMH and testosterone are produced in the testes by different cells and have different effects. Testosterone promotes the development of male genitalia while the binding of AMH to the encoded receptor prevents the development of the mullerian ducts into uterus and Fallopian tubes. Mutations in this gene are associated with persistent Mullerian duct syndrome type II. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2009]
- Synonyms:** AMHR; MISR2; MISRII; MRII
- Note:** This antibody is expected to recognize all reported isoforms (NP_065434.1; NP_001158162.1; NP_001158163.1).
- Protein Families:** Druggable Genome, Protein Kinase, Transmembrane
- Protein Pathways:** Cytokine-cytokine receptor interaction, TGF-beta signaling pathway
- Product images:**



TA348938 (1 ug/ml) staining of MDA-MB-231 lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.