

Product datasheet for **TA348992**

DUSP6 Goat Polyclonal Antibody

Product data:

| | |
|-----------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB: 0.3-1 ug/ml |
| Reactivity: | Human, Mouse, Rat, Pig (Expected from sequence similarity: Dog, Cow) |
| Host: | Goat |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | The immunogen for Anti-DUSP6 / MKP3 Antibody: Peptide with sequence C-PSNQNVYQVDSLQST, from the C Terminus of the protein sequence according to NP_001937.2; NP_073143.2. |
| Formulation: | Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Concentration: | lot specific |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | dual specificity phosphatase 6 |
| Database Link: | NP_001937 Entrez Gene 67603 Mouse Entrez Gene 116663 Rat Entrez Gene 482594 Dog Entrez Gene 1848 Human Q16828 |



[View online »](#)

Background:

The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cytoplasm. Mutations in this gene have been associated with congenital hypogonadotropic hypogonadism. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2014]

Synonyms:

HH19; MKP3; PYST1

Note:


This antibody is expected to recognize both reported isoforms (NP_001937.2; NP_073143.2).

Protein Families:


Druggable Genome, Phosphatase

Protein Pathways:

MAPK signaling pathway

Product images:

TA348992 (0.3 ug/ml) staining of Human (A), (Mouse) and Rat (C) Heart lysates (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



TA348992 (0.3 ug/ml) staining of Pig Heart lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.