

# Product datasheet for TA349271

# **IL20 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: A431 and 231 cells, hela cells and human fetal liver tissue

IHC: 15-50

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Fusion protein of human CDK2

pH7.4 PBS, 0.05% NaN3, 40% Glyceroln Formulation:

Concentration: lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stable for 12 months from date of receipt. Stability:

**Predicted Protein Size:** 34 kDa

Gene Name: interleukin 20 Database Link: NP 061194

Entrez Gene 58181 MouseEntrez Gene 50604 Human

Q9NYY1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



### Background:


The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein kinase is highly similar to the gene products of S. cerevisiae cdc28, and S. pombe cdc2. It is a catalytic subunit of the cyclin-dependent protein kinase complex, whose activity is restricted to the G1-S phase, and essential for cell cycle G1/S phase transition. This protein associates with and regulated by the regulatory subunits of the complex including cyclin A or E, CDK inhibitor p21Cip1 (CDKN1A) and p27Kip1 (CDKN1B). Its activity is also regulated by its protein phosphorylation. Two alternatively spliced variants and multiple transcription initiation sites of this gene have been reported.

Synonyms: IL-20; IL10D; ZCYTO10

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: A431 cells 231 cells hela cells

human fetal liver tissue


Primary antibody: TA349271 (CDK2 Antibody) at

dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA349271 (CDK2 Antibody) at dilution 1/25 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA349271 (CDK2 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)