

## **Product datasheet for TA349373S**

#### rioduct datasileet for 1A349373

# **KIF3A Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse brain and human brain malignant glioma tissue

IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human KIF3A

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 80 kDa

**Gene Name:** kinesin family member 3A

Database Link: NP 008985

Entrez Gene 16568 MouseEntrez Gene 11127 Human

Q9Y496



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### Background:

Kinesin-like protein KIF3A is a protein that in humans is encoded by the KIF3A gene. Members of the heterotrimeric kinesin II family of microtubule associated motors generally contain two different motor subunits from the KIF3 family, which includes KIF3A, B and C. KIF3 isoforms mediate anterograde transport of membrane bound organelles in neurons and melanosomes, transport between the endoplasmic reticulum and the Golgi, and transport of protein complexes within cilia and flagella required for their morphogenesis. KIF3A may influence neurogenesis at the level of embryonic cellular events, where the asymmetry of the genetic control circuit controlling left-right (L-R) axis determination is defined. Loss of KIF3A function in mice photoreceptors causes apoptotic cell death, suggesting that kinesin II mediated transport is required for proper cell fate.

Synonyms: FLA10; KLP-20

**Protein Families:** Druggable Genome

### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: Mouse brain tissue


human brain malignant glioma tissue Primary antibody: [TA349373] (KIF3A Antibody) at

dilution 1/750

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349373] (KIF3A Antibody) at dilution 1/40 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349373] (KIF3A Antibody) at dilution 1/40, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA349373] (KIF3A Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA349373] (KIF3A Antibody) at dilution 1/40, treated with fusion protein. (Original magnification: ×200)