

Product datasheet for TA349449S

HEXIM1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: NIH/3T3 cell, Mouse liver tissue lysates

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human HEXIM1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 41 kDa

Gene Name: hexamethylene bis-acetamide inducible 1

Database Link: NP 006451

Entrez Gene 192231 MouseEntrez Gene 498008 RatEntrez Gene 10614 Human

O94992

Background: Hexamethylene bis-acetamide inducible 1 (HEXIM1) and Hexamethylene bis-acetamide

inducible 2 (HEXIM2) comprise a family of proteins which inhibit positive transcription elongation factor b (P-TEFb) through association with 7SK. P-TEFb is composed of a catalytic subunit, Cdk9, and either Cyclin T1 or T2 as a regulatory subunit. This complex regulates eukaryotic gene expression at the level of elongation. The C-terminal domains of HEXIM proteins interact directly with each other. Via these domains, HEXIM1 and HEXIM2 form stable homo- and hetero-oligomers, which may aid in the formation of the 7SK small nuclear ribonucleic acid particle. Despite their similar functions, HEXIM1 and HEXIM2 exhibit distinct

expression patterns in various established cell lines and human tissues.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: CLP1; EDG1; HIS1; MAQ1

Protein Families: Transcription Factors

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg

Lane 1-2: NIH/3T3 cell Mouse liver tissue lysates

Primary antibody: [TA349449] (HEXIM1 Antibody)

at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at

1/5000 dilution

Exposure time: 1 minute