

Product datasheet for **TA349464S**

AMIGO2 Rabbit Polyclonal Antibody

Product data:

| | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | IHC, WB |
| Recommended Dilution: | WB: 500-2000 WB positive control: SP20 cells and mouse heart tissue IHC: 50-200 Positive control: Human esophagus cancer Predicted cell location: Cytoplasm and Nucleus |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Fusion protein of human AMIGO2 |
| Formulation: | pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol |
| Purification: | Antigen affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 58 kDa |
| Gene Name: | adhesion molecule with Ig-like domain 2 |
| Database Link: | NP_862830 Entrez Gene 105827 MouseEntrez Gene 300186 RatEntrez Gene 347902 Human Q86SJ2 |



[View online »](#)

Background:

The amphoterin-induced gene and ORF (AMIGO) family of proteins consists of AMIGO-1, AMIGO-2 and AMIGO-3. All three members are single pass type I membrane proteins that contain several leucine-rich repeats, one IgG domain, and a transmembrane domain. The AMIGO proteins are specifically expressed on fiber tracts of neuronal tissues and participate in their formation. The AMIGO proteins can form complexes with each other, but can also bind itself. AMIGO-1, also designated Alivin-2, promotes growth and fasciculation of neurites and plays a role in myelination and fasciculation of developing neural axons. In cerebellar neurons, AMIGO-2 (Alivin-1) is crucial for depolarization-dependent survival. Similar to AMIGO-1 and AMIGO-2, AMIGO-3 (Alivin-3) plays a role in homophilic and/or heterophilic cell-cell interaction and signal transduction

Synonyms:

ALI1; AMIGO-2; DEGA

Protein Families:

Druggable Genome, Transmembrane

Product images:

Gel: 6%SDS-PAGE

Lysate: 40 µg


Lane 1-2: SP20 cells

mouse heart tissue


Primary antibody: [TA349464] (AMIGO2 Antibody)
at dilution 1/615

Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution


Exposure time: 2 minutes




Immunohistochemistry of paraffin-embedded
Human esophagus cancer tissue using
[TA349464] (AMIGO2 Antibody) at dilution 1/50
(Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA349464] (AMIGO2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA349464] (AMIGO2 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using [TA349464] (AMIGO2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)