

## **Product datasheet for TA349550S**

## **TLR2 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: K562 cell lysate

IHC: 50-200

Positive control: Human colorectal cancer

Predicted cell location: Cell membrane or Cytoplasm

Reactivity: Human
Host: Rabbit

**Isotype:** IgG

Clonality:

Immunogen: Fusion protein of human TLR2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Polyclonal

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 90 kDa

**Gene Name:** toll like receptor 2

Database Link: NP 003255

Entrez Gene 7097 Human

O60603

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background: The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which


> plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is expressed most abundantly in peripheral blood leukocytes, and mediates host response to Gram-positive bacteria and yeast via stimulation of NF-kappaB.

Synonyms: CD282; TIL4

**Protein Families:** Druggable Genome, Transmembrane **Protein Pathways:** Toll-like receptor signaling pathway

## **Product images:**






Primary antibody: [TA349550] (TLR2 Antibody) at

dilution 1/600

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 20 seconds




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA349550] (TLR2 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA349550] (TLR2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349550] (TLR2 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349550] (TLR2 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)