

## **Product datasheet for TA349565S**

## **SPTLC1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Lovo cells

IHC: 25-100

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human SPTLC1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.


**Predicted Protein Size:** 53 kDa

**Gene Name:** serine palmitoyltransferase long chain base subunit 1

Database Link: NP 847894

Entrez Gene 10558 Human

<u>O15269</u>



9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





**Background:** This gene encodes a member of the class-II pyridoxal-phosphate-dependent


aminotransferase family. The encoded protein is the long chain base subunit 1 of serine palmitoyltransferase. Serine palmitoyltransferase converts L-serine and palmitoyl-CoA to 3-oxosphinganine with pyridoxal 5'-phosphate and is the key enzyme in sphingolipid biosynthesis. Mutations in this gene were identified in patients with hereditary sensory neuropathy type 1. Alternatively spliced variants encoding different isoforms have been identified. Pseudogenes of this gene have been defined on chromosomes 1, 6, 10, and 13.

Synonyms: HSAN1; HSN1; LBC1; LCB1; SPT1; SPTI

Protein Families: Druggable Genome, Transmembrane

**Protein Pathways:** Metabolic pathways, Sphingolipid metabolism

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 µg Lane: Lovo cells


Primary antibody: [TA349565] (SPTLC1 Antibody)

at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349565] (SPTLC1 Antibody) at dilution 1/25 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349565] (SPTLC1 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)