

## **Product datasheet for TA349566S**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## FACL4 (ACSL4) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Hepg2 and hela cells, human fetal kidney and liver tissue

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human ACSL4

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 79 kDa

**Gene Name:** acyl-CoA synthetase long-chain family member 4

Database Link: NP 004449

Entrez Gene 50790 MouseEntrez Gene 113976 RatEntrez Gene 2182 Human

O60488





Background: The protein encoded by this gene is an isozyme of the long-chain fatty-acid-coenzyme A

> ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme preferentially utilizes arachidonate as substrate. The absence of this enzyme may contribute to the mental retardation or Alport syndrome. Alternative splicing of this gene

generates 2 transcript variants.

ACS4; FACL4; LACS4; MRX63; MRX68 Synonyms:

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Adipocytokine signaling pathway, Fatty acid metabolism, Metabolic pathways, PPAR signaling

pathway

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: Hepg2 cells

hela cells

human fetal kidney tissue Human liver tissue


Primary antibody: [TA349566] (ACSL4 Antibody)

at dilution 1/650

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 15 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349566] (ACSL4 Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349566] (ACSL4 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA349566] (ACSL4 Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA349566] (ACSL4 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)