

Product datasheet for TA349617S

Hexokinase II (HK2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: A431, K562, 231, hela and hepG2 cells

IHC: 100-300

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human HK2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 102 kDa

Gene Name: hexokinase 2

Database Link: NP 000180

Entrez Gene 15277 MouseEntrez Gene 25059 RatEntrez Gene 3099 Human

P52789

Background: Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most

glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate

of glycolysis seen in rapidly growing cancer cells.

Synonyms: HKII; HXK2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome


Protein Pathways: Amino sugar and nucleotide sugar metabolism, Fructose and mannose metabolism,

Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Metabolic

pathways, Starch and sucrose metabolism, Type II diabetes mellitus

Product images:





Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-5: A431 cells K562 cells 231 cells hela cells hepG2 cells

Primary antibody: [TA349617] (HK1/HK2

Antibody) at dilution 1/1200


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution


Exposure time: 1 second

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349617] (HK1/HK2 Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349617] (HK1/HK2 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA349617] (HK1/HK2 Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA349617] (HK1/HK2 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)