

## **Product datasheet for TA349622S**

## **ICAM1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Raji and huvec cells

IHC: 50-200

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human ICAM1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 58 kDa

**Gene Name:** intercellular adhesion molecule 1

Database Link: NP 000192

Entrez Gene 3383 Human

P05362

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





**Background:** This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells

and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial

migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation. In case of rhinovirus infection acts as a cellular

receptor for the virus.


Synonyms: BB2; CD54; P3.58

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration, Natural killer cell

mediated cytotoxicity, Viral myocarditis

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-2: Raji cells huvec cells


Primary antibody: [TA349622] (ICAM1 Antibody)

at dilution 1/1150

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 second




Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA349622] (ICAM1 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA349622] (ICAM1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349622] (ICAM1 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA349622] (ICAM1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)