

Product datasheet for TA349640S

BMI1 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-------------------------|---|
| Applications: | WB |
| Recommended Dilution: | WB: 200-1000 WB positive control: 231 and A549 cells, hela cells and mouse brain tissue |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Immunogen: | Fusion protein of human BMI1 |
| Formulation: | pH7.4 PBS, 0.05% NaN3, 40% Glyceroln |
| Purification: | Antigen affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 37 kDa |
| Gene Name: | BMI1 proto-oncogene, polycomb ring finger |
| Database Link: | <u>NP_005171</u> <u>Entrez Gene 12151 MouseEntrez Gene 648 Human</u> <u>P35226</u> |
| Background: | BMI1 polycomb ring finger oncogene, also known as BMI1, is a protein which in humans is encoded by the BMI1gene. Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys- 119', rendering chromatin heritably changed in its expressibility. In the PRC1 complex, it is required to stimulate the E3 ubiquitin-protein ligase activity of RNF2/RING2. |
| Synonyms: | BMI1; FLVI2; PCGF4; RNF51 |
| Protein Families: | Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-4: 231 cells A549 cells hela cells mouse brain tissue Primary antibody: [TA349640] (BMI1 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US