

Product datasheet for TA349650

PIP5K1 beta (PIP5K1B) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PIP5K1B

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: phosphatidylinositol-4-phosphate 5-kinase type 1 beta

Database Link: NP 003549

Entrez Gene 18719 MouseEntrez Gene 309419 RatEntrez Gene 8395 Human

<u>014986</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Phosphatidylinositol-4-phosphate-5-kinase (PIPK) synthesizes phosphatidylinositol-4,5-bisphosphate, which regulates various processes including cell proliferation, survival, membrane trafficking, and cytoskeletal organization. The PIPK family is divided into type I, type II and type III . Each type of the PIPK family phosphorylate distinct substrates and they contain an activation loop, which determines their enzymatic specificity and subcellular targeting . The phosphatidylinositol-4-phosphate-5-kinase type I consists of three members, PIPK I å, \int , and \mathbb{O} , which are characterized by phosphorylating PI4P on the 5-hydroxyl . PIPK I å (designated PIPK I \int in mouse) is expressed in brain tissue . PIPK I \int , designated PIPK I a in mouse, is also called STM7. PIPK I \otimes has two variants produced by alternative splicing which are expressed in lung, brain, and kidneys.


Synonyms: MSS4; STM7

Protein Families: Druggable Genome

Protein Pathways: Endocytosis, Fc gamma R-mediated phagocytosis, Inositol phosphate metabolism, Metabolic


pathways, Phosphatidylinositol signaling system, Regulation of actin cytoskeleton

Product images:




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA349650 (PIP5K1B Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA349650 (PIP5K1B Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using TA349650 (PIP5K1B Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using TA349650 (PIP5K1B Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)