

## **Product datasheet for TA349650S**

## PIP5K1 beta (PIP5K1B) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human PIP5K1B

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

**Gene Name:** phosphatidylinositol-4-phosphate 5-kinase type 1 beta

Database Link: NP 003549

Entrez Gene 18719 MouseEntrez Gene 309419 RatEntrez Gene 8395 Human

O14986

**Background:** Phosphatidylinositol-4-phosphate-5-kinase (PIPK) synthesizes phosphatidylinositol-4,5-

bisphosphate, which regulates various processes including cell proliferation, survival, membrane trafficking, and cytoskeletal organization. The PIPK family is divided into type I, type II and type III . Each type of the PIPK family phosphorylate distinct substrates and they contain an activation loop, which determines their enzymatic specificity and subcellular targeting . The phosphatidylinositol-4-phosphate-5-kinase type I consists of three members, PIPK I å,  $\int$ , and  $\otimes$ , which are characterized by phosphorylating PI4P on the 5-hydroxyl . PIPK I å (designated PIPK I  $\int$  in mouse) is expressed in brain tissue . PIPK I  $\int$ , designated PIPK I a in mouse, is also called STM7. PIPK I  $\otimes$  has two variants produced by alternative splicing which

are expressed in lung, brain, and kidneys.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Synonyms:** MSS4; STM7

**Protein Families:** Druggable Genome


**Protein Pathways:** Endocytosis, Fc gamma R-mediated phagocytosis, Inositol phosphate metabolism, Metabolic

pathways, Phosphatidylinositol signaling system, Regulation of actin cytoskeleton

## **Product images:**




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349650] (PIP5K1B Antibody) at dilution 1/30 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA349650] (PIP5K1B Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human brain tissue using [TA349650] (PIP5K1B Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA349650] (PIP5K1B Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)