

Product datasheet for **TA349662**

ASC1 (TRIP4) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: A431 cells IHC: 25-100 Positive control: Human liver cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human TRIP4
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	66 kDa
Gene Name:	thyroid hormone receptor interactor 4
Database Link:	NP_057297 Entrez Gene 56404 Mouse Entrez Gene 9325 Human Q15650



[View online »](#)

Background:


Activating signal co-integrator-1 (ASC-1), originally identified as TRIP4, is a transcriptional co-activator of nuclear receptors that associates with specific components of the RNA polymerase II complex and binds the basal transcription factors TBP and TFIIA. ASC-1 functions with the transcription integrators SRC-1 and CBP/p300 through its zinc finger motif and is dependent on their ligand-dependent transactivation domain, AF2. Endogenous ASC-1 in HeLa cells is predominantly a nuclear protein. Under conditions of serum starvation, ASC-1 localizes to the cytoplasm.

Synonyms:


ASC-1; ASC1; HsT17391; MDCDC; SMABF1; ZC2HC5

Protein Families:


Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Product images:


Gel: 6%SDS-PAGE
Lysate: 40 µg
Lane: A431 cells
Primary antibody: TA349662 (TRIP4 Antibody) at dilution 1/400
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution
Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA349662 (TRIP4 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA349662 (TRIP4 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: x200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA349662 (TRIP4 Antibody) at dilution 1/50 (Original magnification: x200)



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA349662 (TRIP4 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: x200)