

Product datasheet for TA349859

Cystatin A (CSTA) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 200-1000

WB positive control: Mouse gullet tissue

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human CSTA

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 11 kDa

Gene Name: cystatin A

Database Link: NP 005204

Entrez Gene 1475 Human

P01040

Background: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.

Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: AREI; PSS4; STF1; STFA

Product images:



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Mouse gullet tissue

Primary antibody: TA349859 (CSTA Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 10 minutes