

### **Product datasheet for TA349924**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **EIF3S3 (EIF3H) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human adrenal pheochromocytoma and fetal brain tissue

IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human EIF3H

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 40 kDa

**Gene Name:** eukaryotic translation initiation factor 3 subunit H

Database Link: NP 003747

Entrez Gene 68135 MouseEntrez Gene 299899 RatEntrez Gene 8667 Human

O15372






#### Background:

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

Synonyms:

elF3-gamma; elF3-p40; ElF3S3

# **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg


Lane 1-2: Human adrenal pheochromocytoma and fetal brain tissue

Primary antibody: TA349924 (EIF3H Antibody) at dilution 1/1150

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 40 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA349924 (EIF3H Antibody) at dilution 1/50. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA349924 (EIF3H Antibody) at dilution 1/50. (Original magnification: ×200)