

Product datasheet for TA350036S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIM19 (NDUFA13) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse skeletal muscle and human hepatocellsular carcinoma tissue

IHC: 100-300

Positive control: Human ovarian cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human NDUFA13

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 17 kDa

Gene Name: NADH:ubiquinone oxidoreductase subunit A13

Database Link: NP 057049

Entrez Gene 67184 MouseEntrez Gene 51079 Human

Q9P0J0



Background:


This gene encodes a subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), which functions in the transfer of electrons from NADH to the respiratory chain. The protein is required for complex I assembly and electron transfer activity. The protein binds the signal transducers and activators of transcription 3 (STAT3) transcription factor, and can function as a tumor suppressor. The human protein purified from mitochondria migrates at approximately 16 kDa. Transcripts originating from an upstream promoter and capable of expressing a protein with a longer N-terminus have been found, but their biological validity has not been determined.

Synonyms: B16.6; CDA016; CGI-39; GRIM-19; GRIM19 **Protein Families:** Transcription Factors, Transmembrane

Product images:







Gel: 10%SDS-PAGE Lysate: 40 µg

Lane 1-2: Mouse skeletal muscle human hepatocellular carcinoma tissue Primary antibody: [TA350036] (NDUFA13

Antibody) at dilution 1/350


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution


Exposure time: 5 minutes

Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA350036] (NDUFA13 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA350036] (NDUFA13 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA350036] (NDUFA13 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA350036] (NDUFA13 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)