

## **Product datasheet for TA350049S**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **HMGCL Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse thymus and human ovarian cancer tissue, mouse heart, human

fetal liver, mouse liver tissue and SKOV3 cells

IHC: 50-200

Positive control: Human ovarian cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

Immunogen: Fusion protein of human HMGCL

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 34 kDa

**Gene Name:** 3-hydroxymethyl-3-methylglutaryl-CoA lyase

Database Link: NP 000182

Entrez Gene 15356 MouseEntrez Gene 79238 RatEntrez Gene 3155 Human

P35914

**Background:** The protein encoded by this gene belongs to the HMG-CoA lyase family. It is a mitochondrial

enzyme that catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alternatively spliced transcript variants encoding different isoforms have been found for this

gene.






Synonyms: HL


**Protein Families:** Druggable Genome

**Protein Pathways:** Butanoate metabolism, Metabolic pathways, Synthesis and degradation of ketone bodies,

Valine, leucine and isoleucine degradation


## **Product images:**






Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 5 seconds




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA350049] (HMGCL Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA350049] (HMGCL Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA350049] (HMGCL Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA350049] (HMGCL Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)