

Product datasheet for TA350050S

HMGCR Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse brain tissue lysate

IHC: 50-100

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human HMGCR

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: 3-hydroxy-3-methylglutaryl-CoA reductase

Database Link: NP 000850

Entrez Gene 15357 MouseEntrez Gene 25675 RatEntrez Gene 3156 Human

P04035



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


HMG-CoA reductase is the rate-limiting enzyme for cholesterol synthesis and is regulated via a negative feedback mechanism mediated by sterols and non-sterol metabolites derived from mevalonate, the product of the reaction catalyzed by reductase. Normally in mammalian cells this enzyme is suppressed by cholesterol derived from the internalization and degradation of low density lipoprotein (LDL) via the LDL receptor. Competitive inhibitors of the reductase induce the expression of LDL receptors in the liver, which in turn increases the catabolism of plasma LDL and lowers the plasma concentration of cholesterol, an important determinant of atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms: LDLCQ3

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Metabolic pathways, Terpenoid backbone biosynthesis

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Mouse brain tissue lysate

Primary antibody: [TA350050] (HMGCR Antibody)


at dilution 1/800


Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute









Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA350050] (HMGCR Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)