

## **Product datasheet for TA350057S**

## **HSD17B8 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human testis tissue

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human HSD17B8

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 27 kDa

**Gene Name:** hydroxysteroid (17-beta) dehydrogenase 8

Database Link: NP 055049

Entrez Gene 7923 Human

Q92506



9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





**Background:** In mice, the Ke6 protein is a 17-beta-hydroxysteroid dehydrogenase that can regulate the

concentration of biologically active estrogens and androgens. It is preferentially an oxidative enzyme and inactivates estradiol, testosterone, and dihydrotestosterone. However, the enzyme has some reductive activity and can synthesize estradiol from estrone. The protein encoded by this gene is similar to Ke6 and is a member of the short-chain dehydrogenase superfamily. An alternatively spliced transcript of this gene has been detected, but the full-


length nature of this variant has not been determined.

**Synonyms:** D6S2245E; dJ1033B10.9; FABGL; H2-KE6; HKE6; KE6; RING2; SDR30C1

**Protein Families:** Druggable Genome

**Protein Pathways:** Androgen and estrogen metabolism, Metabolic pathways

## **Product images:**



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane: Human testis tissue


Primary antibody: [TA350057] (HSD17B8

Antibody) at dilution 1/650

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA350057] (HSD17B8 Antibody) at dilution 1/55 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA350057] (HSD17B8 Antibody) at dilution 1/55, treated with fusion protein. (Original magnification: ×200)